


Recommended Outcome Measures along the Total Joint Arthroplasty Continuum



Legend: HOOS = Hip disability and Osteoarthritis Outcome Score; KOOS = Knee injury and Osteoarthritis Outcome Score; LEFS = Lower Extremity Functional Scale; NRPS = Numeric Pain Rating Scale; PROMs = Patient-Reported Outcome Measures; SLS = Single Leg Stance; TUG = Timed-Up-and-Go; VAS = Visual Analogue Scale; 6MWT = 6 Minute Walk Test; 30-CST= 30 Sec Chair Stand Test

The purpose of this document is to provide evidence-based guidance (evidence derived from a survey, chart audit, focus group, Delphi process and the literature) regarding the most appropriate outcome measures (OMs) to use for patients along the continuum of care for Total Hip Arthroplasty and Total Knee Arthroplasty. The selection of how many and which OMs to use is at the discretion of the clinician based upon the unique contexts of the practice setting.

1

Developed by the BC Physical Therapy Total Joint Arthroplasty and Outcome Measures Knowledge Translation Task Force: Dr. Marie Westby, Ronda Field, Maureen Duggan, Dolores Langford, Robyn Laytham, Steven Longstaff, Lauren Welch, Alison Hoens. March 2014; Revised Feb 2015, Revised Feb 2016
A Physical Therapy Knowledge Broker project supported by: UBC Department of Physical Therapy, Physiotherapy Association of BC, Vancouver Coastal Research Institute and Providence Healthcare Research Institute.

REFERENCES

1. Alviar MJ, Olver J, Brand C, Hale T, & Khan F. (2011) Do patient-reported outcome measures used in assessing outcomes rehabilitation after hip and knee arthroplasty capture issues relevant to patients? Results of a systematic review and ICF linking process. *J Rehabil Med*; 43(5):374-81.
2. Alviar MJ, Olver J, Brand C, Tropea J, Hale T, Pirpiris, M, Khan F. (2011). Do patient-reported outcome measures in hip and knee arthroplasty rehabilitation have robust measurement attributes? A systematic review. *J Rehabil Med*; 43:572-83.
3. Collins NJ & Roos EM (2012). Patient-Reported Outcomes for Total Hip and Knee Arthroplasty: Commonly Used Instruments and Attributes of a “Good” Measure. *Clin Geriatr Med*; 28:367–94.
4. Dobson F, Hinman RS, Roos EM, Abbott JG, Stratford P, David AM, Buchbinder R, Snyder-Mackler L, Henrotin Y, Thumboo J, Hansen P, Bennell KL (2013). OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis Cartilage*; 21:1042-52.
5. Hawker GA, Mian S, Kendzerska T, French M (2011). Measures of adult pain. *Arthritis Care Res.* 63(Suppl 11):S240-52.
6. Westby MD, Brittain, A, Backman CL (2013). Expert consensus on best practices for post-acute rehabilitation after total hip and knee arthroplasty: A Canada-US Delphi study. *Arthritis Care Res.* Sep 10. Epub ahead of print.
7. McAuley C, Westby MD, Hoens A, Troughton D, Field R, Duggan M, Reid WD (2014). A survey of physiotherapists' experience using outcome measures in total hip and knee arthroplasty. *Phys Can.* Jan 21. Epub ahead of print.